



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

other hand, *Apios tuberosa*, Mönch, seems to compensate by its tubers for the very frequent abortion of its pods. These plants, with others, *Specularia*, for example, afford an interesting subject for investigation on this point. W. H. L.

7. *Lemna Valdiviana*, Philippi, (L. Torreyi, *Austin*) grows in ponds near Patchogue, Long Island. Its habit is very similar to *L. tib-sulca*, growing under water in masses. I have found only sterile plants. T. F. A.

8. *Nasturtium palustre*, L.—The typical form (smooth, with oblong pods) seems to be quite common about New-York, especially along the Harlem Rail-Road; it is more abundant than the hispid variety. T. F. A.

9. *Arabis laevigata*, DC., grows in abundance on the low, rocky ground just southeast of Yonkers.

10. *Hesperis matronalis* seems thoroughly established, especially in the ditches along the rail-roads. T. F. A.

11. *Bromus s'erilis*, L., is very abundant along the Hudson River R. R., from Manhattanville to Yonkers. It comes very early and lasts but a short time; only dead specimens can usually be obtained after June. T. F. A.

12. The Varian "Sycamore," more properly Plane-tree, which stood on the west side of Broadway, between 26th and 27th streets, was cut down the first week in November. Its diameter was more than four feet near the ground. Just at the street level there was a hollow place on one side, but on the whole it seemed remarkably sound. Some one might give us a history of the stately old giant. Two young observers who counted the rings made out 95, but the roughness of the sections made accuracy difficult.

13. *Polymnia Uvedalia*, L.—Dr. T. F. Allen found a small patch of this plant near the Weehawken Ferry in 1864. It was still to be found there this summer, but in danger of destruction, with the cliffs that sheltered it. Dr. Gray, by some accident, refers this locality to *P. Canadensis*. There is no doubt about its being *P. Uvedalia*.

14. Books.—Dr. Torrey has received "Traité Général de Botanique," by MM. Le Maout and Decaisne, with upwards of 5,500 illustrations by Steinheil and Riocreux.—T. F. A. has Hegelmaier's "Lemnaceen," with 16 plates. He makes three genera, *Wolfia*, (including Griffith's *Grantia*), *Lemna*, and *Spirodela*. L. Torreyi, *Austin*, was less perfectly described by Philippi, 1864, and named *Valdiviana*, from Valdivia, in Chili, where it was first found.—W. H. L. has C. K. Sprengel's "Entdeckte Geheimniss," 1793; often quoted by Darwin.

Subscriptions and communications received by W. H. LEGGETT, 224 E. 10th-street.